

**PROFESSIONAL INFORMATION**
**D 34.12 Multiple Substance Formulation. Complementary Medicine: Health Supplement**

This unregistered medicine has not been evaluated by SAHPRA for its quality, safety or intended use. Health supplements are intended only to complement health or supplement the diet.

**SCHEDULING STATUS: S0**
**1. NAME OF THE MEDICINE**

LOCAL HEALTH MULTIVITAMIN tablets

**2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

Each tablet contains:

Calcium (as Tricalcium Phosphate)	243 mg
Phosphorus (as Tricalcium Phosphate)	125 mg
Magnesium (as Magnesium Oxide)	100 mg
Ascorbic Acid (Vitamin C)	60 mg
Niacin (Vitamin B3)	18 mg
d- $\alpha$ -Tocopherol (Vitamin E)	15 IU / 10 mg TE
Iron (as Electrolytic Iron)	10 mg
Pantothenic Acid (Vitamin B5)	6 mg
Zinc (as Zinc Oxide)	5 mg
Pyridoxine (Vitamin B6)	2 mg
Riboflavin (Vitamin B2)	1,6 mg
Thiamine (Vitamin B1)	1,4 mg
Vitamin A (as Vitamin A Palmitate)	4000 IU / 1200 $\mu$ g
Manganese (as Manganese Sulphate)	1 mg
Copper (as Copper Sulphate 25%)	0,5 mg
Lutein ( <i>Tagetes erecta</i> L. Marigold)	250 $\mu$ g
Folic Acid	195 $\mu$ g
Biotin (Vitamin H)	100 $\mu$ g
Iodine (as Potassium Iodide)	100 $\mu$ g
Selenium (as Selenium Amino Acid Chelate 2%)	60 $\mu$ g
Molybdenum (as Molybdenum Amino Acid Chelate)	50 $\mu$ g
Chromium (as Chromium Amino Acid Chelate 10%)	40 $\mu$ g
Menaquinone (Vitamin K2)	30 $\mu$ g
Cholecalciferol (Vitamin D3)	200 IU / 5 $\mu$ g
Cyanocobalamin (Vitamin B12)	2 $\mu$ g

**Contains sugar:** 88,2 mg isomalt per tablet.

For full list of excipients, see section 6.1.

**3. PHARMACEUTICAL FORM**

Oval, yellow coated tablets.

**4. CLINICAL PARTICULARS**
**4.1 Therapeutic indications**

LOCAL HEALTH MULTIVITAMIN is a daily multivitamin supplement that contains a combination of vitamins, minerals, and antioxidants for the maintenance and support of good health.

**4.2 Posology and method of administration**

Adults 18 years and older: one (1) tablet daily after breakfast.

**4.3 Contraindications**

- Hypersensitivity to any of the active substances or to any of the excipients listed in section 6.1, including members of the Asteraceae/Compositae plant family (ragweed, chrysanthemums, marigolds, daisies, and many other herbs) and iodine.
- Patients with hypercalcaemia, hypercalciuria, or renal impairment.
- Patients with calcium renal calculi or a history of renal calculi.
- Retinoids taken in conjunction with vitamin A containing supplements could have additive toxic effects. Patients should avoid taking vitamin A supplements if they are taking retinoids.
- Patients receiving blood transfusions, parenteral iron therapy, or patients with an anaemia not produced by an iron deficiency, unless iron deficiency is also present, as overdosage may occur.
- Patients with a vitamin B12 deficiency, as folic acid can mask pernicious anaemia by decreasing megaloblastic anaemia. This can prevent appropriate treatment with vitamin B12 and result in neurological damage, such as subacute combined degeneration of the spinal cord.
- Patients with Leber's disease or tobacco amblyopia should not use vitamin B12 as it may cause further degeneration.

**4.4 Special warnings and precautions for use**

- Vitamin C may increase the risk of hyperoxaluria in patients with chronic kidney dysfunction. Large doses (>2 g daily) have been associated with an increased risk of oxalate kidney stones.
- Large amounts of vitamin C can cause haemolysis in individuals with glucose-6-phosphate dehydrogenase (G6PD) deficiency, and can increase the risk of oxalate stone formation in people with a history of oxalate kidney stones. The daily recommended dosage should not be exceeded.
- Patients with iron-storage or iron-absorption diseases such as haemochromatosis or haemoglobinopathies should use iron-containing supplements with caution.
- Patients with gastrointestinal diseases such as inflammatory bowel disease, intestinal strictures or diverticulitis should use iron-containing supplements with caution as iron may cause gastrointestinal irritation and exacerbate these conditions.

**4.5 Interaction with other medicines and other forms of interaction**

- Retinoids taken in conjunction with vitamin A containing supplements could have additive toxic effects. Patients should avoid taking vitamin A supplements if they are taking retinoids.
- Vitamin A taken concomitantly with warfarin may contribute to increased anticoagulant effects.
- Zinc forms various complexes with antibiotics in the gut, affecting the absorption and effectiveness of antibiotics.
- Iron and zinc may interfere with the absorption and effectiveness of penicillamine.
- Thiazide diuretics reduces the urinary excretion of calcium which may increase the risk of hypercalcaemia and milk-alkali syndrome.

- Iron can form toxic complexes with dimercaprol and these should therefore not be administered together.
- Iron and calcium may interfere with the absorption and effectiveness of bisphosphonates.
- Some anticonvulsants (phenytoin, barbiturates, primidone) may reduce the effect of vitamin D by accelerating its metabolism.
- High doses of vitamin D can cause hypercalcaemia. Hypercalcaemia increases the risk of fatal cardiac arrhythmias with digoxin.

**4.6 Fertility, pregnancy and lactation**

Safety in pregnancy and lactation has not been established.

**4.7 Effects on ability to drive and use machines**

No studies on the effects on the ability to drive or use of machinery have been performed. Patients should exercise caution before driving or using machinery until they are reasonably certain that LOCAL HEALTH MULTIVITAMIN does not adversely affect their performance.

**4.8 Undesirable effects**
**Gastrointestinal**

Frequent: gastrointestinal irritation, abdominal pain, nausea, vomiting, diarrhoea, constipation.

**Immune System**

Frequency unknown: hypersensitivity reactions.

**Reporting of suspected adverse reactions**

Reporting suspected adverse reactions after authorization of the medicine is important. It allows continued monitoring of the benefit/risk balance of the medicine. Health care providers are asked to report any suspected adverse reactions to SAHPRA via the "6.04 Adverse Drug Reactions Reporting Form", found online under SAHPRA's publications: <https://www.sahpra.org.za/Publications/Index/8>.

**4.9 Overdose**

In the event of an overdose, undesirable effects as listed in 4.8 can be precipitated or be of increased severity.

Treatment of overdose is symptomatic and supportive.

**5. PHARMACOLOGICAL PROPERTIES**
**5.1 Pharmacodynamic properties**

LOCAL HEALTH MULTIVITAMIN is a daily multivitamin containing a combination of vitamins, minerals and antioxidants for the maintenance of good health.

**5.2 Pharmacokinetic properties**

The active ingredients in this formulation are well known. Pharmacokinetic studies have not been conducted on LOCAL HEALTH MULTIVITAMIN.

**6. PHARMACEUTICAL PARTICULARS**
**6.1 List of excipients**

Magnesium stearate, isomalt, talc, silicon dioxide, polyethylene glycol, yellow coating.

**6.2 Incompatibilities**

Not applicable

**6.3 Shelf Life**

24 months

**6.4 Special precautions for storage**

Store at or below 25 °C.

Store in the original package in order to protect from moisture.

KEEP OUT OF REACH OF CHILDREN.

**6.5 Nature and contents of container**

75 tablets in a white, PET plastic container and white screwcap lid.

**6.6 Special precautions for disposal**

No special requirements.

**7. HOLDER OF CERTIFICATE OF REGISTRATION**

CJ Distribution  
23 Stag Road, Glen Austin, Johannesburg, South Africa  
[careline@cjsa.com](mailto:careline@cjsa.com)  
011 589 2729  
[www.localhealth.com](http://www.localhealth.com)

**8. REGISTRATION NUMBER**

Will be allocated by SAHPRA upon registration.

**9. DATE OF FIRST AUTHORISATION**

Will be allocated by SAHPRA upon registration.

PATIENT INFORMATION LEAFLET

SCHEDULING STATUS

SO

LOCAL HEALTH MULTIVITAMIN (tablets)

Each tablet contains:	
Calcium (as Tricalcium Phosphate)	243 mg
Phosphorus (as Tricalcium Phosphate)	125 mg
Magnesium (as Magnesium Oxide)	100 mg
Ascorbic Acid (Vitamin C)	60 mg
Niacin (Vitamin B3)	18 mg
d-α-Tocopherol (Vitamin E)	15 IU / 10 mg TE
Iron (as Electrolytic Iron)	10 mg
Pantothenic Acid (Vitamin B5)	6 mg
Zinc (as Zinc Oxide)	5 mg
Pyridoxine (Vitamin B6)	2 mg
Riboflavin (Vitamin B2)	1.6 mg
Thiamine (Vitamin B1)	1.4 mg
Vitamin A (as Vitamin A Palmitate)	4000 IU / 1200 µg
Manganese (as Manganese Sulphate)	1 mg
Copper (as Copper Sulphate 25%)	0.5 mg
Lutein ( <i>Tageetes erecta</i> L. Marigold)	250 µg
Folic Acid	195 µg
Biotin (Vitamin H)	100 µg
Iodine (as Potassium Iodide)	100 µg
Selenium (as Selenomethionine 2%)	60 µg
Molybdenum (as Molybdenum Amino Acid Chelate)	50 µg
Chromium (as Chromium Amino Acid Chelate 10%)	40 µg
Menadionine (Vitamin K2)	30 µg
Cholecalciferol (Vitamin D3)	200 IU / 5 µg
Cyanocobalamin (Vitamin B12)	2 µg

Contains sugar: 88.2 mg isomalt per tablet.

Complementary Medicine: Multiple Substance Formulation (Health Supplement)

This unregistered medicine has not been evaluated by SAHPRA for its quality, safety or intended use. Health supplements are intended only to complement health or supplement the diet.

Read all of this leaflet carefully because it contains important information for you.

LOCAL HEALTH MULTIVITAMIN is available without a doctor's prescription, for you to maintain your health. Nevertheless, you still need to use LOCAL HEALTH MULTIVITAMIN carefully to get the best results from it.

- Keep this leaflet. You may need to read it again.
- Do not share LOCAL HEALTH MULTIVITAMIN with any other person.
- Ask your healthcare practitioner or pharmacist if you need more information or advice.

What is in this leaflet?

1. What LOCAL HEALTH MULTIVITAMIN is and what it is used for.

2. What you need to know before you use LOCAL HEALTH MULTIVITAMIN.

3. How to use LOCAL HEALTH MULTIVITAMIN.

4. Possible side effects.

5. How to store LOCAL HEALTH MULTIVITAMIN.

6. Contents of the pack and other information.

1. What LOCAL HEALTH MULTIVITAMIN is and what it is used for

LOCAL HEALTH MULTIVITAMIN is a daily multivitamin supplement that contains a combination of vitamins, minerals, and antioxidants for the maintenance and support of good health.

2. What you need to know before you use LOCAL HEALTH MULTIVITAMIN

Do not use LOCAL HEALTH MULTIVITAMIN:

- If you are allergic or sensitive to any of the ingredients listed in section 6, including members of the Asteraceae/Compositae plant family (ragweed, chrysanthemums, marigolds, daisies, and many other herbs) and iodine.
- If you have hypercalcemia, hypertension, or renal impairment.
- If you have calcium renal calculi, or a history of renal calculi.
- If you are taking retinoids.
- If you are receiving blood transfusions, parenteral iron therapy, or if you have an anaemia not produced by iron deficiency.
- If you have a vitamin B12 deficiency, Leber's disease or tobacco amblyopia.

Warnings and precautions

Consult with a healthcare practitioner prior to using LOCAL HEALTH MULTIVITAMIN, especially if you have a medical condition.

Use LOCAL HEALTH MULTIVITAMIN with caution and consult a healthcare practitioner before use:

- If you have hyperoxaluria or GPD deficiency.
- If you have an iron-storing or iron-absorption disease such as haemochromatosis or haemoglobinopathy.
- If you have an existing gastrointestinal disease such as inflammatory bowel disease, intestinal strictures, or diverticulitis.

Children and adolescents

LOCAL HEALTH MULTIVITAMIN is not recommended for children below the age of 18 years. Consult a healthcare practitioner prior to use.

Other medicines and LOCAL HEALTH MULTIVITAMIN

Always tell your healthcare practitioner if you are taking any other medicine (this includes complementary or traditional medicines). Not all possible interactions with other medicines may be included in this leaflet. Consult with a healthcare practitioner prior to use. LOCAL HEALTH MULTIVITAMIN may interact with the following medications: retinoids, anticoagulants, antibiotics, penicillamine, digoxin, thiazide diuretics, anticonvulsants, dimercaprol, bisphosphonates.

Pregnancy, breastfeeding and fertility

If you are pregnant or breastfeeding, think you may be pregnant or are planning to have a baby, please consult your doctor, pharmacist or other healthcare practitioner for advice before using LOCAL HEALTH MULTIVITAMIN.

Driving and using machinery

No studies on the effects on the ability to drive or use of machinery have been performed. Patients should exercise caution before driving or using machinery until they are reasonably certain that LOCAL HEALTH MULTIVITAMIN does not adversely affect their performance.

3. How to use LOCAL HEALTH MULTIVITAMIN

Do not share medicines prescribed to you with any other person. Do not exceed the daily recommended dosage. Always use LOCAL HEALTH MULTIVITAMIN exactly as described in this leaflet or as your healthcare practitioner has instructed. Check with your healthcare practitioner if you are not sure.

Adults 18 years and older:

One (1) tablet daily after breakfast, or as recommended by a healthcare practitioner.

4. Possible side effects

LOCAL HEALTH MULTIVITAMIN can have side effects. Not all side effects reported for LOCAL HEALTH MULTIVITAMIN are included in this leaflet. Should your general health worsen or if you experience any untoward effects while taking LOCAL HEALTH MULTIVITAMIN, please consult your doctor, pharmacist or other healthcare practitioner for advice. If one of the following happens, stop using LOCAL HEALTH MULTIVITAMIN and tell your doctor or pharmacist immediately, or go to the casualty department at your nearest hospital: swelling of the hands, feet, ankles, face, lips and mouth or throat, which may cause difficulty in swallowing or breathing, rash or itching, fainting. These are very serious side effects. If you have them, you may have had a serious reaction to LOCAL HEALTH MULTIVITAMIN. You may need urgent medical attention or hospitalisation.

• Frequency: abdominal irritation, abdominal pain, nausea, vomiting, diarrhoea, constipation.

• Frequency unknown: hypersensitivity reactions.

If you notice any side effects not mentioned in this leaflet, please inform your doctor or pharmacist.

Reporting of side effects

If you experience side effects, inform your doctor or pharmacist. You can also report side effects to SAHPRA via the "6.04 Adverse Drug Reactions Reporting Form", found online under SAHPRA's publications: <http://www.sahpra.org.za/Publications/Index/8>. By reporting side effects, you can help provide more information on the safety of LOCAL HEALTH MULTIVITAMIN.

5. How to store LOCAL HEALTH MULTIVITAMIN

• Protect from sunlight and store at or below 25 °C.

• Keep out of reach of CHILDREN.

• Do not eat after the expiry date stated on the label.

• Return all unused LOCAL HEALTH MULTIVITAMIN to your pharmacist.

• Do not dispose of unused LOCAL HEALTH MULTIVITAMIN in drains or sewerage systems (e.g. toilets).

6. Contents of the pack and other information

What LOCAL HEALTH MULTIVITAMIN contains

Each tablet contains:

Calcium (as Tricalcium Phosphate)	243 mg
Phosphorus (as Tricalcium Phosphate)	125 mg
Magnesium (as Magnesium Oxide)	100 mg
Ascorbic Acid (Vitamin C)	60 mg
Niacin (Vitamin B3)	18 mg
d-α-Tocopherol (Vitamin E)	15 IU / 10 mg TE
Iron (as Electrolytic Iron)	10 mg
Pantothenic Acid (Vitamin B5)	6 mg
Zinc (as Zinc Oxide)	5 mg
Pyridoxine (Vitamin B6)	2 mg
Riboflavin (Vitamin B2)	1.6 mg
Thiamine (Vitamin B1)	1.4 mg
Vitamin A (as Vitamin A Palmitate)	4000 IU / 1200 µg
Manganese (as Manganese Sulphate)	1 mg
Copper (as Copper Sulphate 25%)	0.5 mg
Lutein ( <i>Tageetes erecta</i> L. Marigold)	250 µg
Folic Acid	195 µg
Biotin (Vitamin H)	100 µg
Iodine (as Potassium Iodide)	100 µg
Selenium (as Selenomethionine 2%)	60 µg
Molybdenum (as Molybdenum Amino Acid Chelate)	50 µg
Chromium (as Chromium Amino Acid Chelate 10%)	40 µg
Menadionine (Vitamin K2)	30 µg
Cholecalciferol (Vitamin D3)	200 IU / 5 µg
Cyanocobalamin (Vitamin B12)	2 µg

Other ingredients: magnesium stearate, isomalt, talc, silicon dioxide, polyethylene glycol and yellow coating.

Contains sugar: 88.2 mg isomalt per tablet.

Presentation and identification of LOCAL HEALTH MULTIVITAMIN

75 oval, yellow coated tablets packed into a white plastic container.

Holder of Certificate of Registration

Manufactured for: CJ Distribution, 23 Stag Road, Glen Austin, Johannesburg, South Africa, 1685

Careline: 011 589 2729; careline@cjdsa.com; www.localhealth.co.za

This pamphlet was last revised in

Will be allocated by SAHPRA upon registration.

Registration number

Will be allocated by SAHPRA upon registration.

PASIËNTINLIGTINGSBLAD

SKEDULERINGSTATUS

SO

LOCAL HEALTH MULTIVITAMIN (tablete)

Elke tablet bevat:	
Kalsium (as Trikalium Fosfaat)	243 mg
Fosfor (as Trikalium Fosfaat)	125 mg
Magnesium (as Magnesiumumoksië)	100 mg
Askorbiensuur (Vitamien C)	60 mg
Niasien (Vitamien B3)	18 mg
d-α-Tokoferol (Vitamien E)	15 IU / 10 mg TE
Iron (as Elektrolytic Iron)	10 mg
Pantotheensuur (Vitamien B5)	6 mg
Zink (as Sinkoksied)	5 mg
Pirdoksien (Vitamien B6)	2 mg
Riboflavin (Vitamien B2)	1.6 mg
Tiamien (Vitamien B1)	1.4 mg
Vitamien A (as Vitamien A Palmitaat)	4000 IU / 1200 µg
Mangan (as Manganusulfat)	1 mg
Koper (as Kopersulfat 25%)	0.5 mg
Lutein ( <i>Tageetes erecta</i> L. Gousblom)	250 µg
Folienuur	195 µg
Biotien (Vitamien H)	100 µg
Jodium (as Kaliumjodid)	100 µg
Selen (as Selen Amonijsourchelaat 2%)	60 µg
Molibdien (as Molibdem Amonijsourchelaat)	50 µg
Chroom (as Chromium Amonijsourchelaat 10%)	40 µg
Menadionien (Vitamien K2)	30 µg
Cholekalisiferol (Vitamien D3)	200 IU / 5 µg
Cyanocobalamin (Vitamien B12)	2 µg

Bevat suiker: 88.2 mg isomalt per tablet.

Komplementêre Medisyne: Meervoerde Bestanddele Formulasie (Gesondheidsaanvulling)

Hierdie onregistreerde medisyne is nie deur SAHPRA vir gebruik, veiligheid of beoogde gebruik geëvalueer nie. Gesondheidsaanvullings dien slegs as aanvulling tot gesondheid of diete.

Lees hierdie inligtingsblad noukeurig daar want dit bevat inligting wat belangrik is vir u.

LOCAL HEALTH MULTIVITAMIN is beskikbaar sonder 'n dokter se voorskrif, om u gesondheid te handhaaf. Nietemin moet LOCAL HEALTH MULTIVITAMIN versigtig gebruik word ten einde die beste resultate te verkry.

- Hou hierdie inligtingsblad. Dit kan nodig wees dat u dit weer lees.
- Moenie LOCAL HEALTH MULTIVITAMIN met enige ander persoon deel nie.
- Vra u gesondheidsoorgeskryf van apoteker indien u meer inligting nodig het.

Wat is in hierdie inligtingsblad

1. Wat LOCAL HEALTH MULTIVITAMIN is en waarvoor dit gebruik word.
2. Wat u moet wet voordat u LOCAL HEALTH MULTIVITAMIN gebruik.
3. Hoe om LOCAL HEALTH MULTIVITAMIN te gebruik.
4. Moontlike nieue-effekte.
5. Berging van LOCAL HEALTH MULTIVITAMIN.
6. Inhoud van die verpakking en ander inligting.

1. Wat LOCAL HEALTH MULTIVITAMIN is en waaroor dit gebruik word

LOCAL HEALTH MULTIVITAMIN is 'n daagliks multivitamineaanvulling wat 'n kombinasie van vitamine, mineraale en antioksidente bevat vir die handhawing en ondersteuning van gode gesondheid.

2. Wat u moet voordat u LOCAL HEALTH MULTIVITAMIN gebruik

Moenie LOCAL HEALTH MULTIVITAMIN neem nie:

- Indien u alergie of sensitiwiteit teen enige van die bestanddele gelys in afdeling 6, insluitende lede van die Asteraceae / Compositae plantfamilie (dissel, krisante, groenteblomme, madeliefies en ander kruisblomme) en jodium.
- Indien u hiperkalemie, hiperkalorie of mineralenbeperking het.
- Indien u kaliumniveuse het, of 'n geskeleidens van nierstene het.
- Indien u retroinjekte gebruik.
- Indien u bloedopretappings of parenterale ysterterapie ontvang, of as 'n anemie het wat nie deur ysteretekort veroorsaak word nie.
- Indien u in vitamine B12-teker. Leber se siekte of tabak-amblyopie het.

Weerskuings en voorwaardes van gebruik

Raadpleeg 'n gesondheidsoorgeskryf voordat u LOCAL HEALTH MULTIVITAMIN gebruik, veral as u 'n mediese toestand het.

Gebruik LOCAL HEALTH MULTIVITAMIN met omsigtlike en raadpleeg 'n gesondheidsoorgeskryf vir gebruik:

- Indien u in hiperkalemie of GPD-teker het.
- Indien u ysteropberings- of ysterabsorpse-siekte het, soos hemochromatose of hemoglobinoopathie.
- Indien u in gastritis en intestinale siekte het, soos inflammatoriese dermsiekte, intestinale vernouings of divertikula.

Kinders en adolescentes

LOCAL HEALTH MULTIVITAMIN word nie aanbeveel vir kinders jonger as 18 jaar nie. Raadpleeg 'n gesondheidsoorgeskryf vir gebruik.

Die naam van LOCAL HEALTH MULTIVITAMIN en ander medisyne

Lig altyd 'n gesondheidsoorgeskryf in indien u enige ander medisyne neem (dit sluit aanvullende of tradisionele medisyne in). Nie alle moontlike interaksies met ander medisyne weet in hierdie pamphlet ingesluit nie. Raadpleeg 'n gesondheidsoorgeskryf vir gebruik. LOCAL HEALTH MULTIVITAMIN kan interaksies hê met die volgende medisyne: retinoids, anticoagulants, antibiotics, penicillamine, digoxin, thiazide diuretics, anticonvulsants, dimercaprol, bisfosfonates.

Swangerskap, borsvoeding en vrugbaarheid

Raadpleeg 'n gesondheidsoorgeskryf voordat u LOCAL HEALTH MULTIVITAMIN gebruik, veral as u geswanger bent.

Geen studies is uitgevoer op die effekte op die vernoë om te bestuur of die gebruik van masjinerie nie. Pasiënte moet versigtig wees voordat hulle bestuur of masjinerie gebruik, tottdat hulle redelik seker is dat LOCAL HEALTH MULTIVITAMIN nie hul werkverrigting nadig beïnvloed nie.

3. Hoe om LOCAL HEALTH MULTIVITAMIN te gebruik

Moenie medisyne wat u vreesgelyk is met enige ander persoon deel nie. Moenie die aanbeveelde daglikse dosis oorskry nie. Gebruik LOCAL HEALTH MULTIVITAMIN presies soos aanbeveel in hierdie inligtingsblad, of soos aanbeveel deur 'n gesondheidsoorgeskryf. Raadpleeg 'n gesondheidsoorgeskryf indien u nie seker is nie.

Volwassenes 18 jaar en ouer:

Neem een (1) tablet daagliks na ontby, of soos aanbeveel deur 'n gesondheidsoorgeskryf.

4. Moontlike nieue-effekte

LOCAL HEALTH MULTIVITAMIN kan nieue-effekte hê. Nie alle nieue-effekte vir LOCAL HEALTH MULTIVITAMIN is ingesluit in hierdie inligtingsblad nie. Indien u algemeen gesondheid verwerger, of as u enige ongewone reaksieservaardigheid van die gebruik van LOCAL HEALTH MULTIVITAMIN, raadpleeg 'n gesondheidsoorgeskryf vir advies. As een van die volgende gebeur, hou op met die gebruik met LOCAL HEALTH MULTIVITAMIN en raadpleeg 'n gesondheidsoorgeskryf vir advies om te stuur of hou op met enige ongewone reaksie: swelling of houteblaas, gal en fluitse. Hierdie is baie ernstige nieue-effekte. Indien jy dit het, mag jy 'n ernstige allergiese reaksie op LOCAL HEALTH MULTIVITAMIN gehad het. Jaag dringende mediese aandag van hospitaalsieheid nadig hê.

- Diukwels: gastritis en intestinale iritisie, bukgyn, narakeid, brak, diarree, hardlywrigheid.
- Diukwels onbekend: hypersensitiviteitsreakties.

Lig altyd 'n gesondheidsoorgeskryf in indien u enige nieue-effekte opmerk wat nie in hierdie inligtingsblad genoem word nie.

Rapportering van ongewone reaksies

Praat met 'n dokter of apoteker as u ongewone reaksieservaardigheid van die gebruik van LOCAL HEALTH MULTIVITAMIN, wat gevind word deur SAHPRA aangevolg via die "6.04 Adverse Drug Reactions Reporting Form", wat gevind word deur SAHPRA as publikasie: <http://www.sahpra.org.za/Publications/Index/8>. Deur nieue-effekte aan te meld, kan u help om meer inligting te verskaf vir die veiligheid van LOCAL HEALTH MULTIVITAMIN.

5. Bewaring van LOCAL HEALTH MULTIVITAMIN

• Bakskerm teen varsel en bly op onder 25 °C.

• HOU DIETIE-DEREK VAN KINDEREN.

• Moenie gebruik na die verwadelut op die etiket nie.

• Neem alle ongebruikte LOCAL HEALTH MULTIVITAMIN na u apoteker terug.

• Moenie ongebruikte LOCAL HEALTH MULTIVITAMIN in afvoerlypte of röolsisteme (bv. toilette) weggooi nie.

6. Inhoud van die verpakking en ander inligting

Wat LOCAL HEALTH MULTIVITAMIN bevat

Elke tablet bevat:	





<tbl\_r cells="2" ix="5" maxcspan="1" maxrspan="1" usedcols="